

ABOS Trauma Recertification Examination Blueprint

ABOS Trauma Recertification Examination Blueprint Table of Contents

Pelvis/Acetabulum	8-12%
Upper Extremity	18-22%
Lower Extremity	29-33%
Polytrauma	8-12%
General/Basic Science	19-21%
Geriatrics	6-10%

ABOS Trauma Recertification Examination Blueprint

Pelvis/Acetabulum	8-12%
Pelvis (initial)	1-3%
Pelvis (definitive)	1-3%
Acetabulum	1-4%

Upper Extremity	18-22%
Shoulder-scapula	0.5-1%
Shoulder-clavicle	1-2%
Shoulder-proximal humerus	1-4%
Humerus shaft	1-2%
Elbow-distal humerus	1-3%
Elbow-proximal/radius/ulna/dislocations	1-3%
Elbow-pediatric supracondylar humerus	0.5-1%
Forearm-radius/ulna	0.5-1%
Distal radius	1-3%

Lower Extremity	29-33%
Hip-femoral head (young patient), femoral neck, dislocations	1-2%
Geriatric hip-proximal femur	2-5%
Femur, shaft and subtrochanteric	1-3%
Adolescent femur	0.5-1%
Distal femur	1-3%
Knee-patella and extensor mechanism	0.5-1%
Knee-dislocations	0.5-1%
Tibia, proximal	1-3%
Tibial shaft	1-2%
Tibia, distal (pilon)	1-2%
Ankle fracture/dislocations	1-4%
Foot, talus/calcaneus	1-2%
Mid-foot/forefoot	1-2%

Polytrauma	8-12%
Acute respiratory distress syndrome/ Systemic inflammatory response syndrome/ Fat embolism syndrome	1-2%
Orthopaedic management of the polytrauma patient	1-4%
Initial assessment of polytrauma patient-ABCs	1-2%
Initial assessment of polytrauma patient-head	0.5-1%
Initial assessment of polytrauma patient-spine	0.5-1%

General/Basic Science	19-21%
Outcomes (e.g., PTSD, mental health, measures)	1-2%
Biomechanics of implants	0.5-1%
Thromboembolic/anticoagulation/bleeding control	1-2%
Mangled extremity/amputation/prosthetics/orthotics	1-3%
Malunion/nonunion	1-2%
Infection/osteomyelitis/antibiotics	1-2%
Soft tissue/open fractures	1-4%
Bone grafts/bone graft substitutes	1-2%
Compartment syndrome	0.5-1%
Pain management/opioids	0.5-1%
Host risk factors (smoking, diabetes, obesity, malnourishment)	0.5-1%

Geriatrics	6-10%
Osteoporosis, diagnosis and management	1-3%
Pathologic fractures/atypical	1-2%
Periprosthetic fractures	1-2%
Perioperative management of geriatric fracture patients	1-3%